

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

CD2 monoclonal antibody, clone MEM-65 (Biotin)

Catalog Number: MAB4556

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against native CD2.

Clone Name: MEM-65

Immunogen: Native purified CD2 from human

peripheral T cells.

Host: Mouse

Theoretical MW (kDa): 50

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: This antibody recognizes an unique epitope of CD2, a 50 KDa glycoprotein present on the human peripheral blood T-lymphocytes and NK cells; also

expressed by all thymocytes.

Form: Liquid

Conjugation: Biotin

Isotype: IgG1

Recommend Usage: Flow Cytometry (1:1000)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS, pH 7.2 (0.09% sodium azide)

Storage Instruction: Store at 4°C. Do not freeze. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 914

Gene Symbol: CD2

Gene Alias: SRBC, T11

References:

1. CD2 deficiency partially prevents small bowel inflammation and improves parasite control in murine Toxoplasma gondii infection. Pawlowski NN, Struck D, Grollich K, Kuhl AA, Zeitz M, Liesenfeld O, Hoffmann JC. World J Gastroenterol. 2007 Aug 21;13(31):4207-13.

2. Mechanisms of Cellular Avidity Regulation in CD2-CD58-Mediated T Cell Adhesion. Zhu DM, Dustin ML, Cairo CW, Thatte HS, Golan DE. ACS Chem Biol. 2006 Nov 21;1(10):649-58.

3. Adhesive bond dynamics in contacts between T lymphocytes and glass-supported planar bilayers reconstituted with the immunoglobulin-related adhesion molecule CD58. Dustin ML. J Biol Chem. 1997 Jun 20;272(25):15782-8.