

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

ITGAL monoclonal antibody, clone MEM-25 (Biotin)

Catalog Number: MAB4563

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against native ITGAL.

Clone Name: MEM-25

Immunogen: Native purified ITGAL from leukocytes of

patient suffering from LGL-type leukaemia.

Host: Mouse

Theoretical MW (kDa): 170-180

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: This antibody reacts with CD11a (alpha-subunit of human LFA-1), a 170-180 KDa type I transmembrane glycoprotein expressed on B and T lymphocytes, monocytes, macrophages, neutrophils,

basophils and eosinophils.

Form: Liquid

Conjugation: Biotin

Isotype: IgG1

Recommend Usage: Flow Cytometry (1:120)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS, pH 7.2 (0.09% sodium azide)

Storage Instruction: Store at 4 °C. Do not freeze. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 3683

Gene Symbol: ITGAL

Gene Alias: CD11A, LFA-1, LFA1A

Gene Summary: ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

References:

- 1. Roles played by lymphocyte function-associated antigen-1 in the regulation of lymphocytic cholinergic activity. Fujii T, Takada-Takatori Y, Kawashima K. Life Sci. 2007 May 30;80(24-25):2320-4. Epub 2007 Jan 17.
- 2. Membrane-proximal signaling events in beta-2 integrin activation. Kellersch B, Kolanus W. Results Probl Cell Differ. 2006;43:245-57.
- 3. Monoclonal antibodies against human leucocyte antigens. IV. Antibodies against subunits of the LFA-1 (CD11a/CD18) leucocyte-adhesion glycoprotein. Bazil V, Stefanova I, Hilgert I, Kristofova H, Vanek S, Horejsi V. Folia Biol (Praha). 1990;36(1):41-50.